

## **Blue Waters Annual Report 2017 – Research Summary Submission**

### **Title: High Resolution Numerical Simulation of Oscillatory Flow and Sediment Transport through Aquatic Vegetation**

**PI:** Rafael O. Tinoco Lopez, Department of Civil and Environmental Engineering, University of Illinois at Urbana-Champaign

**Co-PI:** Paul Fischer, Department of Computer Science and Department of MechSE, University of Illinois at Urbana-Champaign

#### **Collaborator(s):**

Som Dutta, Department of Computer Science, University of Illinois at Urbana-Champaign

Pallav Ranjan, Department of Civil and Environmental Engineering, University of Illinois at Urbana-Champaign

**Corresponding author:** Rafael O. Tinoco Lopez, [tinoco@illinois.edu](mailto:tinoco@illinois.edu)

#### **Executive Summary**

Aquatic vegetation provides a wide range of services to the ecosystem: improving water-quality through nutrient uptake and oxygen production, providing flood buffering and coastal protection services, and regulating erosion and deposition patterns; thus playing a paramount role in habitat creation and promotion of biodiversity. While vegetation-flow interactions have been extensively studied for unidirectional flows, relatively much less is known for oscillatory conditions. The current study is geared towards increasing our understanding of the interactions between vegetation, flow and sediment under oscillatory flows. Direct Numerical Simulations (DNS) and Large Eddy Simulations (LES) through different arrays of idealized vegetation, represented as cylinders, are conducted using the higher-order spectral element based computational fluid dynamics (CFD) solver Nek5000. Different arrangements and numbers of cylinders have been simulated in 2D and 3D, with the largest simulation having ~ 296 million computational points, using up to 32,768 MPI ranks.

#### **Key Challenges**

Our study is geared to identify fundamental physical processes driving interactions between flow, vegetation and sediment, conducting numerical simulations at unprecedented scales, based on previous and ongoing experiments at the Ven-Te-Chow Hydrosystems Laboratory at UIUC. Flow through random and staggered arrays of cylinders is investigated to understand the effect of spatial heterogeneity of the vegetation on the flow. The study primarily focus on oscillatory flow, though a few cases of unidirectional flow will be conducted for comparison purposes. Coupling the experimental and numerical study will yield further understanding of sediment dynamics under the influence of vegetation.

A better understanding of the vegetation-flow dynamics, will not only advance fundamental knowledge of physical processes, but also guide design efforts for scour protection and artificial wetlands. Part of the study is to characterize drag-coefficients of the vegetation array, improving accuracy of reduced-order models of flow through vegetation. The study will also identify the ideal quantity and locations to place instrumentation on experiments through large arrays of cylinders.

### **Why it Matters.**

Seagrasses are commonly referred to as “ecosystem engineers” due to its ability to modify and stabilize its environment [1]. They are a fundamental component of near-shore ecosystems, providing a wide-range of services [2], ranging from increase in water-quality through nutrient uptake and oxygen production, creation of habitats through spatial heterogeneity of the flow velocity, to dampening erosion on coastal. Past studies have focused mostly in unidirectional flows, relying strongly on experimental approaches only [3,4], with limited applications to oscillatory conditions. Such experiments provide ambient conditions closer to nature, although their measurements often lack the spatial and temporal resolution required to fathom the fundamental physical processes in detail. On the other hand, most numerical studies to date have primarily used CFD models based on temporal averaging of the Navier-Stokes equations (RANS), which approximates the turbulence in the system rather than accurately calculating it, and a few LES studies had to settle for a relatively small number of vegetation elements.

### **Why Blue Waters?**

The study pushes the limit of the scale at which high-resolution simulations are used to study complex multi-phase flow in environmental fluid mechanics, requiring computational resources with sustained computing power at an unprecedented scale. The number of computation points required to model the whole domain is near 1.2 billion. While such simulations are still tractable on a petascale platform like Blue Waters, the computational cost is abhorrently high; resulting in reduction of number of cases one could run, thus constraining the insights a broader range of parameters could yield. To increase variable space, a wide range of conditions are first simulated in 2D for the whole domain (~ 4 million computational points), to get an overview of the effect of different parameters like Reynolds number, Vegetation Density, Period and Amplitude of the Oscillatory flow, and spatial heterogeneity. Once the effects of different parameters are well understood, 3D simulations will be conducted for partial domains, big enough to accurately capture the general dynamics, but within manageable computational costs. Without access to petascale HPC like Blue Waters, completing the study within a realistic timeframe would be impossible. In addition, since visualization of a phenomenon is an effective way to understand and explain its mechanics, we will work with Blue Waters project staff to create animations of the phenomenon using data from the simulations.

## Accomplishments

High-resolution Large Eddy Simulations (LES) and Direct Numerical Simulations (DNS) of the flow at different configurations of the idealized vegetation are conducted using the open-source, spectral element based higher-order incompressible Navier-Stokes solver Nek5000 [5]. The Spectral Element Method (SEM) combines the accuracy of spectral methods and the flexibility of Finite Elements Method (FEM) [7]. In the planned simulations with sediment transport, sediment would be modeled as Lagrangian particles using a novel semi-implicit time stepping scheme developed to simulate polydisperse sediment accurately [8].

2D simulations have been conducted for the full domain on different configurations. Two cases having the same vegetation density and Reynolds number, but different array configuration, are presented herein. The velocity magnitude, along with the pressure field, are shown in Fig. 1. For the staggered case, in contrast with the random array, a vortex being shed from a cylinder is impeded by the ones behind it. This is evident in the pressure plots, where more and larger low-pressure areas, indicating the low-pressure core of rotating vortices, appear in the random case. For comparison, the random configuration was subjected to unidirectional flow at the same Reynolds number (Fig. 2). High-flow zones near the walls are noticed, resulting in stronger vortices being shed from near-wall cylinders. Compared with the oscillatory flow case, more high-speed regions are also found among cylinders on the unidirectional case.




Figure 1: Instantaneous velocity magnitude (a,c) and hydrodynamic pressure (b,d) for oscillatory flow at Reynolds number  $\sim 10000$ . For staggered (a,b) and random (c,d) configuration of cylinders the flow has been captured while accelerating from left to right. For the staggered case, the vortex being shed from a cylinder is impeded by the ones behind it, which is not the case for the random case. This is also obvious from the hydrodynamics pressure plots, where the random case present more abundant low-pressure regions, characteristic of the low-pressure core of rotating vortices.




Figure 2: Instantaneous velocity magnitude (a) and hydrodynamic pressure (b) for a random array of cylinders with unidirectional flow of Reynolds number  $\sim 10000$ , flowing from left to right. Notice the high-flow zones near the walls, which results in stronger vortices being shed from near-wall cylinders.




Figure 3: Results from 3D simulation of turbulent flow through a random arrangement of cylinders. About 296 million computational points are used. Instantaneous velocity magnitude at different elevations. Planes at (a) 0.5 %, (b) 1 %, (c) 10 %, (d) 50 %, (e) 75 %, and (f) 95 % water depth. Notice the length of the vortex being shed increases with distance from the bottom.



Results from 3D simulations of turbulent flow through a random arrangement of cylinders have also been conducted (Fig. 3). A quarter of the full domain is simulated, with ~296 million computational points. Fields of instantaneous velocity magnitude at different elevations show that the length of the vortex being shed increases with distance from the bottom. This simulation, being one of the largest high-resolution eddy resolved hydrodynamic simulation in this field, is to provide yet unseen details on the physical processes involved.

## References

- [1] Koch, E. W. (2001). Beyond light: physical, geological, and geochemical parameters as possible submersed aquatic vegetation habitat requirements. *Estuaries*, 24(1), 1-17.
- [2] Nepf H.M. Hydrodynamics of vegetated channels. *Journal of Hydraulic Research*. 2012 Jun 1;50(3):262-79.
- [3] Nepf, H. (2012b). Flow and transport in regions with aquatic vegetation. *Ann. Rev. Fluid Mech.* 44, 123–142.
- [4] Tinoco, R., Coco, G., 2016. A laboratory study on sediment resuspension within arrays of rigid cylinders. *Adv. Water Res.* 92, 1–9.
- [4] Dutta, S., et al., Application of computational fluid dynamic modelling to improve flow and grit transport in Terence J. O'brien water reclamation plant, Chicago, Illinois, *J. Hydr. Res.*, 52:6 (2014), pp. 759-774.
- [5] Fischer, P.F., J.W. Lottes, and S.G. Kerkemeier, Nek 5000 webpage, (2008), <http://nek5000.mcs.anl.gov>
- [6] Deville, M., P. Fischer, and E. Mund, High-Order Methods for Incompressible Fluid Flow, *Cambridge, England: Cambridge University Press*, (2002).
- [7] S. Dutta, P. Fischer, M.H. Garcia, A novel semi-implicit Lagrangian particle tracking model for efficient computation of particle transport at low Stokes number, *under preparation for Journal of Scientific Computing, Springer*.

## List of publications, data sets associated with this work.

- {1} S. Dutta, P. Fischer, M.H. Garcia, A novel semi-implicit Lagrangian particle tracking model for efficient computation of particle transport at low Stokes number, under preparation for Journal of Scientific Computing, Springer.
- {2} S. Dutta, P. Fischer, M.H. Garcia, Large Eddy Simulation (LES) of flow and bedload transport at an idealized 90-degree diversion: insight into Bulle-Effect, under preparation for Journal of Scientific Computing, Springer.

{3} Dutta, S., Ranjan, P., Mittal, K., Fischer, P., & Tinoco, R.O. "2D High-resolution numerical study of emergent vegetation: characterizing drag on unidirectional flows", under preparation for Env. Fluid Mech.

{4} Dutta, S., Mittal, K., Ranjan, P., Fischer, P., & Tinoco, R.O. "2D High-resolution numerical study of emergent vegetation: The impact of vegetation layout under oscillatory flows", under preparation for Env. Fluid Mech.